

Wouter Arrazola de Oñate, MD Public Health



BIJZONDER MODEL VOOR OPVANG VAN TUBERCULOSEPATIËNTEN

BELTA-TBnet is een project dat met financiële steun van de federale overheid werd opgericht. Het verzekert de terugbetaling van de kosten verbonden aan de tuberculosebehandeling die niet door de mutualiteit, het OCMW of een andere sociale instelling ten laste genomen worden.

[Lees meer...](#)

15. Institut Georges Born. **Wenduyne s/Mer**
La façade principale.

Gesticht Georges Born
De voorgevel

1897

BELTA 
Belgian Lung and
Tuberculosis Association



**BELGISCH NATIONAAL WERK
TOT BESTRIJDING DER TUBERCULOSE**

Instelling van Openbaar Nut

Vereeniging zonder winstgevend doel



TE ELSENE


STANDREGELLEN

European disease – not a ‘tropical’ disease



land grab termed the “scramble for Africa” in the late 19th century, placing vast portions of the African continent under European control in the form of colonies. Our findings thus suggest an intimate relationship between European colonial expansion and the spread of L4 TB on the African continent. The main form of






fedasil



alias.brussels







COMMENT · Volume 403, Issue 10443, P2467-2469, June 08, 2024

[Download Full Issue](#)

Tuberculosis: a biosocial problem that requires biosocial solutions

Anurag Bhargava^a · Madhavi Bhargava^b · Madhukar Pai^c 






Same building

Patients
TB-nurses
Physicians



Social cheques
Food - transport
Housing - shelter



Outreach -Include

– BEST PRACTICES IN PREVENTION, CONTROL AND CARE FOR DRUG- RESISTANT TUBERCULOSIS

A resource for the continued implementation of the Consolidated Action Plan to Prevent and Combat Multidrug- and Extensively Drug-Resistant Tuberculosis in the WHO European Region, 2011–2015




Submitted by: Maryse Wanlin, Wouter Arrazola de Oñate and Guido Groenen – Belgian Lung and Tuberculosis Association




BELGIUM

BELTA-TBnet, an example of best practices in the context of MDR-TB management




BELTA 
 Belgian Lung and
 Tuberculosis Association


2023




REDEN TEN LASTE NEMING BELTA-TBNET IN 2024



Évolution du nombre absolu des personnes prises en charge par BELTA-TBnet selon l'indication



Uitgaven in EUR



MDR-TB regimens:

BPaLM
 BPAL

> 50,000 euro/pt

Functioneringskosten Onderzoeken/raadplegingen
 Geneesmiddelen Totaaluitgaven

BELTA - MDR-TB Expert Group

pres: prof Natalie Lorent



Concilium: PH, pediatric, pneumo, infectio, lab...

- Clinical
- Lab/WGS – Vanessa Mattys & team (NRL) + Leen Rigouts (ITM)


Individualised treatment




BPaL - BPaLM

Access to drugs

Guidelines









Team work





Ministry of Health
Republic of Poland



POLAND25.EU



Moving Forward on HIV, viral hepatitis, TB and STIs

**Progress towards reaching the
Sustainable Development Goal 3.3
targets related to HIV, tuberculosis,
viral hepatitis and sexually
transmitted infections in the EU/EEA**


Belgium. BELTA-DOT: intensified and individualized case management for TB patients to improve compliance and treatment outcomes in Belgium

Submitted by: Wouter Arrazola de Oñate^{1,2} | Lilas Weber^{1,3} | Guido Groenen^{1,4} | Lies Geysens^{1,2,5} | Kristien Janssens^{1,2} | Maryse Wanlin^{1,3} | Sandrina Schol^{1,2}

Compendium of good practices
in the implementation of the
Tuberculosis Action Plan for the
WHO European Region 2016–2020



Intensified individualised Case Management + integrated care network



Antwerpen Brussel

Projectmedewerker Outreach - Community Health

- uit Afghanistan aangekomen minderjarigen
- jongvolwassenen uit Eritrea-Ethiopië (Hoorn van Afrika)

(cultural competence) (cultural humility)

“Racisme sensitieve zorg” - dr Ama Kissi (UGent)

6.395 bereikt met gezondheidsinformatie
over TB

805 RX'en

TB-detectiegraad: 497 per 100.000

